



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

thalloid development in ferns than in the *Ophioglossaceae* and a difference in the evolution of the spores. He would, therefore, place the *Ophioglossaceae* in an order of equivalent value with the *Filices*, but in advance of them in a system of classification, inasmuch as they are in some respects more highly differentiated than the latter. —The *Library Journal* for November contains an article by Prof. Ezra Abbot, of Harvard University, showing to what extent many of our standard works of reference continue to perpetuate the old and erroneous view as to the papyrus (*Papyrus antiquorum*.) For instance in "Adam's Roman Antiquities" we read that this plant was "about ten cubits high, and had several coats or skins above one another, like an onion, &c." In Smith's "Dictionary of Greek and Roman Antiquities," under the article *Liber*, the writer says: "The *papyrus-tree* grows in swamps, &c.," and that "paper was prepared from the *thin coats or pellicles which surround the plant*." Liddell and Scott's Greek Lexicon defines βίβλος as "the *inner bark* of the papyrus." A similar account is given in the Lexicon of Jacobitz and Seiler, Pape, and Rost and Palm's edition of Passow under βίβλος and πάπυρος; so also in many encyclopaedias. *e. g.*, the "Encyclopaedia Britannica," and others. This common error of speaking of the *papyrus* as if it were an exogenous plant (and even a *tree*!) has originated from ignorance or forgetfulness of the elements of botany, and the consequent misinterpretation of the passage in Pliny (*Hist. Nat.* xiii. 11-13, al. 21-27), which is our chief source of information about the ancient manufacture of paper from this plant. One of the words Pliny uses to describe the thin strips into which the *cellular substance of the stem* was sliced in making the paper is *philyra*, which strictly denotes the inner bark of the Linden tree (*Tilia*), also used as a writing material. Hence the papyrus has been conceived of by the eminent authorities above cited as an exogen, with its inner and outer bark!

W. R. G.

§ 300. *Epigaea repens*, L.—I found specimens of this plant in full bloom at Princes Bay, S. I., on Saturday, Mch. 1st. I do not know that it has ever been found earlier in this locality.

A. H.

§ 301. *Anychia dichotoma*, Mchx., **not dichotomous**.—I do not know whether attention has already been called to the fact, that the specific name of this plant is really a misnomer, if we take the term, "dichotomy," in its strict scientific signification.

If we examine younger specimens of our *Anychia*, we invariably find every axis terminated by a *flower*, with a branch on each side from *lateral* buds *below* the apex. Hence this is a plain case of cymose, not of dichotomous ramification. In older specimens, say toward the end of July or in August, when many of these terminal flowers have fallen off, the main stem and many branches appear bifurcated. But I need not repeat that this cannot be called dichotomy, which only occurs when some axis, *at its very apex*, is "cut into two" branches, which may again be divided in the same manner, and so on.

Now, although I commonly agree with those who believe in the